



TITLE:

Patterns of morphological variation in enamel-dentin junction and outer enamel surface of human molars.

AUTHOR(S):

Morita, Wataru; Yano, Wataru; Nagaoka, Tomohito; Abe, Mikiko; Ohshima, Hayato; Nakatsukasa, Masato

CITATION:

Morita, Wataru ...[et al]. Patterns of morphological variation in enamel-dentin junction and outer enamel surface of human molars.. *Journal of anatomy* 2014, 224(6): 669-680

ISSUE DATE:

2014-04-01

URL:

<http://hdl.handle.net/2433/198842>

RIGHT:

This is the peer reviewed version of the following article: Morita, W., Yano, W., Nagaoka, T., Abe, M., Ohshima, H. and Nakatsukasa, M. (2014), Patterns of morphological variation in enamel–dentin junction and outer enamel surface of human molars. *Journal of Anatomy*, 224: 669–680, which has been published in final form at <http://dx.doi.org/10.1111/joa.12180>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ください。

1 **Original Paper**

2 Title: Patterns of morphological variation in enamel-dentin junction and outer-enamel surface of human
3 molars

4 Authors: Wataru Morita*, Wataru Yano, Tomohito Nagaoka, Mikiko Abe, Hayato Ohshima, Masato
5 Nakatsukasa

6 Affiliations:

7 Wataru Morita*, Laboratory of Physical Anthropology, Department of Zoology, Graduate School of
8 Science, Kyoto University, Kitashirakawaoiwakecho, Sakyo-ku, Kyoto 606-8502, Japan, Tel.:
9 +81-75-753-4094, E-mail: morita@anthro.zool.kyoto-u.ac.jp

10 Wataru Yano, Department of Oral Anatomy, Division of Oral Structure, Function and Development,
11 Asahi University School of Dentistry, Mizuho, Gifu 501-0296, Japan, Tel.: +81-58-329-1404, E-mail:
12 yano@dent.asahi-u.ac.jp

13 Tomohito Nagaoka, Department of Anatomy, St. Marianna University School of Medicine, Kawasaki,
14 Kanagawa 216-8511, Japan, Tel.: +81-44-977-8111, E-mail: nagaoka@marianna-u.ac.jp

15 Mikiko Abe, Department of Anatomy and Cell Biology, Graduate School of Medicine, Osaka City
16 University, Osaka 545-8585, Japan, Tel.: +81-6-96645-3706, E-mail: mikiko@med.osaka-cu.ac.jp

17 Hayato Ohshima, Division of Anatomy and Cell Biology of the Hard Tissue, Department of Tissue
18 Regeneration and Reconstruction, Niigata University Graduate School of Medical and Dental Sciences,
19 2-5274 Gakkocho-dori, Chuo-ku, Niigata 951-8514, Japan, Tel.: +81-25-227-2812, E-mail:
20 histoman@dent.niigata-u.ac.jp

21 Masato Nakatsukasa, Laboratory of Physical Anthropology, Department of Zoology, Graduate School

22 of Science, Kyoto University, Kitashirakawaoiwakecho, Sakyo-ku, Kyoto 606-8502, Japan, Tel.:

23 +81-75-753-4094, E-mail: nakatsuk@anthro.zool.kyoto-u.ac.jp

Abstract

Tooth crown patterning is governed by the growth and folding of the inner enamel epithelium (IEE) and the following enamel deposition forms outer enamel surface (OES). We hypothesized that overall dental crown shape and covariation structure is determined by processes that configurate shape at enamel-dentin junction (EDJ), the developmental vestige of IEE, and tested this hypothesis by comparing patterns of morphological variation between EDJ and OES in human maxillary permanent first molar (UM1) and second deciduous molar (um2). Using geometric morphometric methods, we described morphological variation and covariation between EDJ and OES, and evaluated the strength of two components of phenotypic variability: canalization and morphological integration, in addition to the relevant evolutionary flexibility, i.e., the ability to respond to selective pressure. The strength of covariation between EDJ and OES was greater in um2 than UM1, and the way that multiple traits covary between EDJ and OES were different between these teeth. The variability analyses showed that EDJ had less shape variation and a higher level of morphological integration than OES, which indicated that canalization and morphological integration acted as developmental constraints. These tendencies were greater in UM1 than um2. On the other hand, EDJ and OES had the comparable level of evolvability in these teeth. Amelogenesis could play a significant role in tooth shape and covariation structure, and its influence was not constant among teeth, which may be responsible for the differences in the rate and/or period of enamel formation.

Key Words: developmental constraints, geometric morphometrics, morphological variability, evolvability, odontogenesis

Introduction

Dental morphological characteristics such as cusps, accessory cusps, and ridges on the occlusal surface have been used extensively in the studies of hominoid evolution and phylogenetic relationships (Miller, 1918; Simons and Pilbeam, 1972; Dean, 2000; Pilbrow, 2006; Matsumura et al., 2011). Tooth crown morphology is determined through two developmental processes (Avishai et al., 2004; Skinner and Gunz, 2010; Smith et al., 2011). The first process is the growth and folding of the inner enamel epithelium (IEE) during the bell stage. This morphogenesis (= tooth crown patterning) is governed by interactions between the IEE and underlying mesenchymal tissues. The final configuration of the IEE is preserved as the enamel-dentin junction (EDJ). The second process is biomineralization by the enamel-forming ameloblasts and dentin-forming odontoblasts. Ameloblasts are derived from the IEE cells and odontoblasts from the dental papilla cells. Enamel formation starts at the cusp tips, and proceeds apically to complete the outer-enamel surface (OES).

Recent micro-CT dental analyses have revealed that crown morphological traits of the completed EDJ are modified or masked through the process of enamel deposition (Skinner et al., 2009, 2010; Ortiz et al., 2012), and that the extent of modification varies depending, in part, if not totally, on the enamel thickness (Ortiz et al., 2012). This raises a concern about whether or not shared derived features and homoplastic features of similarity at the OES can be properly discriminated (Hunter and Jernvall, 1995; Collard and Wood, 2000; Finarelli and Clyde, 2004). Additionally, by examining enamel thickness variation and its heritability in pedigreed baboon molars, Hlusko et al. (2004) showed that enamel thickness could change rapidly under moderate or low selective pressure over evolutionarily short periods, increasing the potential for homoplasy. Although the OES morphology is directly related

to dental functions such as occlusion and feeding and thus is a direct target of natural selection, the morphology of EDJ has been considered to be more conservative evolutionally and a more reliable representation of the phenotype for estimating phylogenetic relationships (Kraus, 1952; Korenhof, 1960; Smith et al., 1997; Sasaki and Kanazawa, 1999; Smith et al., 2000; Olejniczak et al., 2007).

So far researchers have explored to which extent enamel formation influences on the crown morphology by comparing EDJ with OES (Kraus, 1952; Nager, 1960; Korenhof, 1960, 1961; Sakai and Hanamura, 1971; Skinner et al., 2008, 2009; Ortiz et al., 2012). However, these studies mainly have focused on discrete dental traits. Although a few studies tried to evaluate general morphological difference between EDJ and OES quantitatively by using intercuspal distance (Smith et al., 1997) or surface complexity (Skinner et al., 2010), complex dental crown topography of EDJ and OES has not been clarified in detail. Examining morphological variation and covariation between EDJ and OES enables us to understand the effects of morphological change caused by enamel formation.

Additionally, given the different developmental backgrounds between the EDJ and OES, it is likely that the patterns of phenotypic variability differ between these structures. Phenotypic variability is defined as the tendency or potential of an organism to vary (Wagner and Altenberg, 1996; Wagner et al., 1997; Willmore et al., 2007). Therefore, it determines the potential range or distribution of morphological variation, and ultimately affects the tempo and mode of evolutionary change. The recent literature about phenotypic variability has paid the greatest attention to canalization and morphological integration (Wagner and Altenberg, 1996; Hallgrímsson et al., 2002; Willmore et al., 2007; Hallgrímsson et al., 2009). Canalization is generally considered a property of an organism that limits phenotypic variation by buffering developmental processes against both environmental and genetic

perturbations (Wagner et al., 1997; Willmore et al., 2007). Morphological integration refers to the tendency for different characters to covary as a result of common underlying developmental factors (Hallgrímsson et al., 2002), which constrains the production of phenotypic variation (Wagner and Altenberg, 1996; Chernoff and Magwene, 1999). Canalization and morphological integration are interrelated and can act as developmental constraints (Alberch, 1982; Maynard Smith et al., 1985; Hallgrímsson et al., 2002). Since the morphological integration framework is directly connected to the rate and direction of evolutionary change (Cheverud, 1996; Wagner and Altenberg, 1996), some studies have focused on quantification of the intervening effect of morphological integration on evolutionary trajectory (Lande, 1979; Lande and Arnold, 1983). The resultant data have led to recent studies that evaluated evolvability (the ability of a population or species to respond to selection: Hansen, 2003) using the simulation of evolutionary responses to selection (Marroig et al., 2009; Villmoare et al., 2011; Lewton, 2012; Grabowski, 2013). The relationships and interactions among developmental processes, variability and variation, mediated by the feedback loop of natural selection, are critically involved in evolutionary change (Willmore et al., 2007). Comparison of the pattern of variability between EDJ and OES helps to infer how the production of morphological variation is regulated in each of these components.

In this study, we explore the relationship between the crown morphology and odontogenesis through quantitative analyses of the EDJ and OES morphology. We hypothesized that overall dental crown shape and covariation structure are determined by processes that configure shape at the EDJ. If this hypothesis is rejected, a significant role of enamel formation for patterning of crown morphological variation must be presumed. To test this hypothesis, we described morphological variation and

108 covariation between EDJ and OES and revealed how much variation in the OES shape is explained by
109 the EDJ shape variation. Consequently, we evaluated the strength of two components of phenotypic
110 variability: canalization and morphological integration, in addition to the relevant evolutionary
111 flexibility.

112 *Canalization*: if EDJ shows larger variation, it means that more variable morphology is created during
113 the early phase of the tooth development, and subsequent enamel formation acts as stabilizing
114 developmental process that buffers the deviation from mean shape. On the other hand, if OES shows
115 larger variation, it indicates that enamel formation brings about not only homogeneous enamel
116 distribution above the EDJ after the morphogenesis, but also some modification of the OES associated
117 with the increased variation.

118 *Morphological integration*: if either during morphogenesis or the enamel formation process, some
119 developmental factors produce higher morphological integration of one of these structures (whether
120 EDJ or OES). Combined with the result regarding canalization, this analysis will help to determine what
121 factors play important roles in generating or reducing morphological variance.

122 *Evolutionary flexibility*: in relation to the above two components of phenotypic variability, we
123 specifically compared how the developmental constraints exert influence on the ability of the response
124 to selection in EDJ versus in OES.

125 This study focused on EDJ and OES shape variation of maxillary permanent first molar
126 (UM1) and second deciduous molar (um2). Although UM1 and um2 share similar patterns of occlusal
127 morphology that are elaborated through the same developmental processes, their developmental timing,
128 speed and period are different (Nanci, 2013). The differences between UM1 and um2 will provide a

better understanding of the relationship between odontogenesis and crown morphological variability.

Materials and Methods

The samples used in this study comprised fully formed but unworn UM1 and um2 crowns obtained from archaeological sites in Japan. The total sample (57 UM1 and 48 um2) consisted of samples from the Jomon (14500-300 BC; n=8 and 5), Medieval (13-15C AD; n=13 and 8), and Edo (17-19C AD; n=36 and 35) periods. Although the total sample was from a mixture of populations from different periods and regions, the aim of this study was to investigate differences and patterns of variability produced by a common tooth formation process of the Holocene human, and mixing these samples does not violate the objective of this study. In order to maximize sample size, no discrimination between right and left teeth was made, but only a single tooth was used from each individual. All specimens were regarded as left side. Right molar images were transformed into the mirror image using ImageJ software (NIH, USA). Sex was unknown for most of the samples, since they were taken from juvenile individuals.

Each specimen was μ CT scanned (ScanXmateA080S, Comscantecno, Japan) with a pixel size and slice interval of 31–32 μ m (80 kV, 125 μ A). To facilitate tissue segmentation, the image stack for each tooth was filtered using a median filter followed by a kuwahara filter, and enamel and dentin tissues were segmented by the seed region growing method in ImageJ. Triangular mesh models of the 3D EDJ and OES of each specimen were reconstructed using Analyze 6.0 (Mayo Clinic, USA) with the marching cube method. Subsequent procedures were done using the software Rapidform 2004 (INUS Technology, Korea).

We treated the EDJ and the OES as biologically corresponding structures in order to compare variability between them directly, and digitized (semi)landmarks on both of them in the same way as follows. We digitized main cusp tips (paracone, protocone, metacone, and hypocone) at the OES and the dentin horn tips at the EDJ, and the lowest points on the ridges at both the OES and the EDJ, connecting the two cusps as landmarks. Each ridge on both the OES and the EDJ was divided into eight sections by the cusp tips and the lowest points, respectively. For each section, a given number of semi-landmarks was digitized equi-distantly, as illustrated in Figure 1. The number of semi-landmarks on the EDJ and the OES were determined to satisfy two criteria, namely, that each corresponding section in the EDJ and the OES had the same number of (semi)landmarks, and that the contributions of the section between the (semi)landmarks to the curve were approximately equal to each other (Skinner et al., 2009; Skinner and Gunz, 2010). The dataset consisted of four configurations (UM1EDJ, UM1OES, um2EDJ and um2OES), and each of them had a total of 8 landmarks and 48 semi-landmarks.

Semi-landmarks are not considered to be homologous landmarks unless they are slid (Bookstein, 1997). The minimum bending energy algorithm (Bookstein, 1997; Gunz et al., 2005) was adopted. This data processing was performed by W. Y. using MATHEMATICA 8 (www.wolfram.com). Each homologous set of landmarks was converted to shape coordinates by Generalized Procrustes Analysis (GPA; Rohlf and Slice, 1990), which was performed using MorphoJ version 1.05d (Klingenberg, 2011).

Morphometric analysis

Covariation between EDJ and OES was analyzed using 2B-PLS. This method compares two morphological data sets by using a singular value decomposition of the cross-covariation matrix, finds new pairs of axes that account for the maximum amount of covariance between both data sets and visualizes the main associated morphological changes. The RV coefficient was used to evaluate the strength of multivariate correlations between data sets. This coefficient is a multivariate analogue of the squared correlation coefficient (Escoufier, 1973; Klingenberg, 2008). The significances of both the correlation between the scores for each pair of PLS axes and RV coefficient were evaluated by means of resampling tests with 1000 random permutations. These procedures were carried out with MorphoJ software (Klingenberg, 2011).

A principal component analysis of Procrustes shape coordinates was used to extract main patterns of morphological variation across EDJ and OES in both UM1 and um2. Using first few PC scores of EDJ and OES, we performed a regression analysis between these two structures to test whether shape variation of OES can be predicted by that of EDJ.

The difference in multivariate morphological change vector from EDJ to OES between UM1 and um2 was assessed by calculating the length and direction of shape change using a residual randomization procedure outlined in Collyer and Adams (2007). The length of a vector describes the overall amount of morphological change and the direction of a vector describes the way in which multiple traits covary. Observed vector lengths and directions were compared with 999 random permutations plus the observed value to assess significance.

Variability analysis

Among-individual phenotypic variation is the most common measurement of canalization.

Canalization is generally inferred from a reduction of the observed phenotypic variance. Here we quantified both size and shape variance within each of the four configurations. For size, Centroid size (CS) of each configuration was calculated. Coefficient of variation (CV) of the LogCS was used to compare size variation, and tested as suggested by Sokal and Braumann (1980). For comparison of shape variability among configurations, the square root of the sum of the squared distances between Procrustes transformed coordinates of each cusp and its landmark mean configuration was used as the measure of shape variation. To test whether there was a significant difference of variability between the EDJ and the OES within the same tooth class, a nonparametric Kruskal-Wallis test and multiple-comparison test were performed.

To compare the overall strength of morphological integration, we followed Wagner (1984) in using the variance of the eigenvalues for the variance-covariance matrix as the measure of integration. This measure of integration captures whether shape variance can be explained by a small number of principal components, or whether variance is more evenly distributed across principal components. The former case would be considered more integrated and the latter less integrated. Variance of eigenvalues (VE) was compared between the EDJ and the OES within the same tooth using bootstrap resampling methods (Manly, 1997). For each of the EDJ and the OES, the original data matrix was bootstrapped 1000 times, a variance-covariance matrix was derived from each bootstrap sample, and VE was calculated from each of the 1000 variance-covariance matrices. For each of the 1000 VE replicates, the difference between the EDJ and the OES was calculated. This created a distribution of differences in VE replicates that was then zero-centered. Each of the zero-centered differences was then compared to

the observed difference in VE between the EDJ and the OES. The two-tailed P value was calculated as the number of times the difference from the zero-centered distribution was equal to or greater than the observed difference, divided by the number of bootstrap replicates (Manly, 1997).

The ability of EDJ and OES morphology to respond to selection was evaluated using mean flexibility (f) (Marroig et al. 2009), which is derived from Lande's (1979) multivariate selection equation:

$$\Delta \mathbf{z} = \mathbf{G} \boldsymbol{\beta}$$

where \mathbf{G} is the genetic covariance matrix, $\boldsymbol{\beta}$ is a selection vector, and $\Delta \mathbf{z}$ is the response vector. Here the phenotypic covariance matrix \mathbf{P} is substituted for \mathbf{G} because previous studies established structural similarity between them (e.g., Cheverud, 1996; Porto et al., 2009). The covariance matrix for each of EDJ and OES was subjected to 1,000 randomly generated selection vectors and the angle between the selection and response vectors was calculated for each time. The mean cosine of angles in 1000 repeats is called the mean flexibility (Marroig et al., 2009), which describes the degree to which the response and selection vectors are aligned in multivariate space. Response and selection vectors that are parallel (i.e., when the cosine of the angle between them is 1.0) indicate a structure that is more responsive to selection, i.e., more evolvable. A larger angle between the response and selection vectors is indicative of less evolvability. In general, high levels of evolvability measures, such as evolutionary flexibility, tend to be associated with low levels of integration measures (e.g., VE). Pairwise comparisons of evolutionary flexibility between EDJ and OES within the same tooth class were performed as described for VE; the distribution of vector correlations obtained from the covariance matrix and 1,000 random selection vectors for EDJ and OES were compared using the difference of means test and accompanied

by a two-tailed P value. All statistical analyses were performed using R version 2.13.1 (R Development Core Team, 2011).

Results

Morphometric analysis

Covariation between EDJ and OES is higher in um2 ($RV=0.914$; $P<0.001$) than in UM1 ($RV=0.794$; $P<0.001$). 2B-PLS analysis in UM1 revealed that the first axis explained 49.43% of total shape covariance and that corresponding shape change mainly involves the contraction of buccal side and expansion of distolingual cusp (hypocone) for both EDJ and OES (Table 1; Fig. 2a). The second axis also revealed that EDJ and OES showed similar shape change that contraction of mesiobuccal cusp (paracone) and contraction of distal side (Fig. 2b). In um2, the first singular axis of correspondence to the comparison of EDJ and OES revealed a correlated reduction of mesiolingual-distobuccally and expansion of mesiobuccal-distolingually (Fig. 2c). The second axis also revealed significant shape change of reduction of mesial cusps and reduction of distal cusps for both EDJ and OES (Fig. 2d).

In the PCA, the first two principal components account for 34.85% of the total variation (Figure 3a; Table 2). Positive scores of PC1 are associated with relatively high and sharp cusp tips and lingually located hypocone. Its negative values correspond to relatively-gentle and inner located cusp tips with deep intercuspal grooves. Positive PC2 scores are associated with mesial expansion and contraction of protocone and negative ones with mesial contraction with lingually expanded protocone. PC1 corresponds to the distinction between EDJ and OES, whereas PC2 separates between UM1 and um2. Figure 3b and 3c illustrates the regressions of first two PCs for EDJ and OES in both teeth. The

adjusted R- squared value is lower in UM1 than that in um2 for both PC1 (0.249 vs.0.700) and PC2 (0.842 vs. 0.907), which indicated that the OES shape variation is better predicted by EDJ shape variation in um2 than in UM1.

The tooth specific morphological change vectors between EDJ and OES were not statistically different in length ($\Delta D=0.004$; $P=0.27$). However, the angle between these vectors was significantly greater than expected by chance ($\theta =27.62^\circ$; $P<0.001$; Fig. 3a).

Variability analysis

Canalization

The coefficients of variation of the LogCS for each configuration (UM1EDJ, UM1OES, um2EDJ and um2OES) was not significantly different from each other, although OES tended to be more variable than EDJ in both the UM1 and um2 tooth classes (Figure 4a). On the other hand, shape variability was significantly different among these configurations, and pair-wise tests showed that only in UM1 was there a significant difference in shape variability between EDJ and OES (Figure 4b).

Morphological integration

The variance of the eigenvalue (VE) was significantly greater for EDJ than for OES in UM1, but not in um2 (Figure 4c). The greater VEs for EDJ were seen in both UM1 and um2, indicating that EDJ was more integrated than OES.

Evolutionary flexibility

The mean cosine between the selection vector and the response vector for OES tended to be greater than that for EDJ, but a significant difference was not detected between them in either tooth class (Figure 4d). This meant that there was no difference in the extent to which EDJ and OES would be influenced by the selection vector.

Discussion

Both UM1 and um2 showed significantly correlated shape changes between EDJ and OES corresponding to singular axes. Enamel formation does not alter the basic morphology of the dentine horn and EDJ ridges and corresponding features (cusp tips and ridges) on OES. Our results accord with previous studies that dental traits seen in EDJ can be observed at OES (Korenhof, 1961, 1982; Nager, 1960; Sakai and Hanamura, 1973; Corruccini, 1998; Sasaki and Kanazawa, 1999; Skinner et al., 2008; 2009), which supports the major role of the EDJ in their origin and degree of dental crown traits. However, this does not necessarily mean that tooth shape and covariation structure are predetermined by processes that configure tooth shape at EDJ. Comparisons between um2 and UM1 revealed different influences of enamel formation on the OES morphology. In um2, OES shape variation is better predicted from EDJ shape variation. Thus, multivariate covariation between EDJ and OES is higher compared to UM1. This result suggests that morphological change caused by enamel formation is more stable and less vulnerable to random perturbations in um2. This could be attributed to the difference in the enamel thickness (Grine, 2005), the rate of enamel formation (Shellis, 1984) and/or period of enamel formation (Liversidge and Molleson, 2004). While the amount of overall morphological change induced by enamel formation does not differ between UM1 and um2, the direction of change described

297 by traits covariation marks a significant difference. Given the different period of formation between
298 UM1 and um2 (Nanci, 2013), it may be expected that they show resembling directions of
299 morphological change with different amounts of morphological change. However, the result is converse,
300 suggesting a complex nature of crown enamel formation. For example, Grine (2005) noted that the
301 difference in enamel thickness between the paracone tip and the protocone tip was greater in um2 rather
302 than in UM1. The difference in patterns of enamel distribution between UM1 and um2 might affect the
303 way of covariation between EDJ and OES. Thus, enamel formation has a significant effect on *patterns*
304 of morphological change, probably according to tooth-specific developmental parameter, though it does
305 not cause a drastic change in morphology during odontogenesis.

306 The lack of significant difference in size variation between EDJ and OES in both tooth
307 classes examined here suggests that the strength of canalization on size is almost constant throughout
308 the processes of morphogenesis and the subsequent enamel formation period. A recent developmental
309 study revealed that molar crown sizes were regulated by intrinsic factors from mesenchymal tissues (Cai
310 et al., 2007) and adjacent molars during development (Kavanagh et al., 2007). Several dental metrics
311 studies confirmed that tooth crown size was less variable than intercuspal distance and/or cusp size owing
312 to stronger genetic control (Townsend et al., 2003; Harris and Dihn, 2006), which would be also
313 supported by experimental evidence that cusp density (intercuspal distances) was likely to be polygenic
314 (Harjunmaa et al., 2012). The present analysis of EDJ and OES at the dentin horns/cusp tips and ridges
315 provided the insight about intercuspal distances that their size variation might not be altered largely by
316 enamel formation. Additionally, the spatial relationship with the surrounding tissues, including the
317 maxillary bone and/or other tooth germs, and the available space for tooth growth (Boughner, 2011)

may be involved in the canalization of crown size during odontogenesis. The extent of the deviation from mean size in EDJ and OES were not significantly different, and therefore both EDJ size differences and OES size differences among groups being compared can be used as a reliable measure of phylogenetic relatedness.

In the case of UM1, shape variation of OES was greater than that of EDJ. This result suggests that canalization of crown shape may be weakened during the process of enamel formation. Kraus and Jordan (1965) argued that early stages of tooth development were mediated by genes that are more evolutionarily stable than those associated with calcification. Hlusko's (2004) simulation model indicated that enamel thickness could change rapidly under appropriate selective pressure. The present result obtained at the cusp tips and ridges is in accord with these studies and implies that shape (e.g., intercusp topological relationship) variation is more susceptible to modifications resulting from enamel formation than size variation, which might be likely to cause homoplasy that would confuse phylogenetic reconstructions (note here "size" refers to the centroid size of the cuspal tips and ridges and not commonly used crown size proxies like maximum mesiodistal x buccolingual dimensions).

The result of VE analysis showed that EDJ was more integrated than OES in UM1, although the same was not supported statistically in um2. Molar crown morphogenesis is a morphodynamic process in which inductive events and morphogenetic processes act at the same time, and is regulated by interactions between the epithelial and underlying mesenchymal tissues. Cusp initiation and patterning in tooth germ is an iterative process that repeatedly utilizes the same set of genes and signaling pathways, which would lead to higher morphological integration in EDJ. On the other hand, the pattern of enamel formation is the end product of a sequence proceeding from ameloblast differentiation from the IEE

cells, to secretion of enamel proteins including amelogenins and enamelines, and finally organization of the enamel crystallites into enamel rods or prisms (Boyde, 1964, 1989). Topological developmental parameters, such as the rate and the duration of enamel apposition and/or ameloblast extension and termination (Simmer et al., 2010), might impact the OES formation, which could lead to weaker morphological integration in OES.

It is predicted that stronger integration between traits acts as a limitation on producing phenotypic variation (Wagner and Altenberg, 1996). The results of the canalization and morphological integration analyses presented here are consistent with this prediction, i.e., the more strongly integrated EDJ shows smaller variability. The set of genes expressed during morphogenesis of the tooth are also used in different organs, including hair, pancreas, mammary gland, salivary gland, thymus, vibrissae, and others (Fincham et al., 2000; Jernvall and Jung, 2000). Mutations in coding region that alter the function or activity of proteins are likely to have widespread and potentially many negative effects on development and fitness, and may thus be under considerable constraint (Carroll, 2008). Size and shape of EDJ are thus more likely to be stabilized in order to reduce the risks of negative pleiotropic side effects. The high level of integration in EDJ can be regarded as a relatively rigorous developmental constraint during odontogenesis. Meanwhile, the set of genes that contribute to enamel formation, such as amelogenin, enamelin, ameloblastin, and enamelysin genes, is highly specialized, and can easily modify the OES morphology during the enamel formation process. Morphological change of the OES, which has less developmental constraint, can easily be brought about by neutral evolution by non-natural selective genetic factors such as random genetic drift.

The observed pattern of morphological integration and the results of evolutionary flexibility

analyses presented here are not consistent with those of previous studies, in which low levels of integration accompanied high levels of evolvability (Marroig et al., 2009; Porto et al., 2009; Lewton, 2012). The developmental constraints due to canalization and morphological integration act more strongly on the shape of EDJ than on that of OES in UM1, while there is no significant difference in the evolutionary flexibility between EDJ and OES. This may result from the relatively integrated covariance structure of each cusp (for both EDJ and OES). Since the secondary enamel knot that functions as a signaling center and regulates cusp formation at the future cusp tip acts as a “developmental module” (Jernvall and Jung, 2000), it can directly affect the covariance structure of EDJ, and indirectly affect that of the overlying OES. In the case of the human tooth, if the crown covariance structure is divided into individual cusp units, this patterning cascade mode of cusp development facilitates the ability to respond to selective challenges (Jernvall and Jung, 2000), and enables the maintenance of a certain level of evolvability at EDJ despite existence of developmental constraints. The comparable level of evolutionary flexibility between EDJ and OES suggests that both of them can be utilized as an equally effective proxy for inferring phylogenetic relationships that would result from selective pressure.

Overall, the difference of each measurement (canalization, morphological integration and evolutionary flexibility) between the EDJ and OES in the present study was greater in UM1 than in um2. The process of enamel formation is more likely to influence crown morphological variability and evolvability in UM1 than in um2, which can be explained by the duration and/or thickness of enamel formation. Compared to UM1, the enamel deposition period of um2 is shorter and the enamel is thinner (Nanci, 2013). Therefore enamel formation may exert less influence on shape change in um2, which

may be related to the conservation of primitive morphology, as discussed in previous studies (Dahlberg, 1945; Butler, 1956, 1971; Suzuki and Sakai, 1973; Saunders and Mayhall, 1982). Since not only morphology but also variability would be likely to differ between EDJ and OES, a tooth crown that has a longer period of enamel formation and/or thicker enamel would require careful evaluation for phylogenetic studies.

This study compared patterns of canalization, morphological integration, and evolutionary flexibility between the EDJ and the OES in UM1 and um2 in order to explore their possible effects on phylogenetic reconstructions. Our results suggest that a tooth crown that has thicker enamel and/or a longer period of enamel formation can be more variable in shape at the OES, where similarity can be due to homoplasy. Recent advances in imaging techniques have made it possible to approach the details of developmental trajectories reflected in the teeth of fossil species (Avishai et al., 2004; Smith et al., 2011). Understanding the morphological variability and evolvability produced by the developmental process is an important step in validating phylogenetic hypotheses based on the OES morphology alone.

Concluding Remarks

Both morphometric and variability analyses indicate that tooth shape and covariation structure is not only determined by processes that contribute to tooth shape at the EDJ, but also amelogenesis can play a significant role in them. The influence of enamel formation on morphological variation and patterns of variability is not constant among teeth, which may be responsible for the differences in the rate and/or period of enamel formation.

402 **Acknowledgements**

403 The authors thank Y. Kunitatsu, D. Shimizu, N. Morimoto, and other members of the
404 Laboratory of Physical Anthropology, Kyoto University, for helpful discussions and comments, P.
405 Gunz and P. Mitteroecker for permission to use software routines they developed, and K. Hirata, K.
406 Shimatani, and K. Miyazawa for access to specimens under their care. We are also grateful to Editor
407 and to two anonymous reviewers for their constructive criticism. This study was supported in part by a
408 JSPS Research Fellowship (11J00940), and by Grants for Excellent Graduate Schools, MEXT, Japan.
409 The authors declare no potential conflicts of interest with respect to the authorship and/or publication of
410 this article.

411

412 **Author contributions**

413 W.M. and W.Y. designed the research and performed the analysis. W.M., T.N., and M.A.
414 collected the data. W.M., H.O., and M.N. wrote the manuscript.

415 **References**

- 416 Alberch P (1982) Developmental constraints in evolutionary processes. In: *Development in evolution*
417 (ed. Bonner JT), pp. 313–332. Berlin and New York: Springer-Verlag.
- 418 Avishai G, Müller R, Gabet Y, Bab I, Zilberman U, Smith P (2004) New approach to quantifying
419 developmental variation in the dentition using serial microtomographic imaging. *Microsc Res Tech* 65,
420 263–299.
- 421 Bookstein FL (1997) Landmark methods for forms without landmarks: morphometrics of group
422 differences in outline shape. *Med Image Anal* 1, 225–243.
- 423 Boughner JC (2011) Making space for permanent molars in growing baboon (*Papio anubis*) and great
424 ape (*Pan paniscus* and *P. troglodytes*) mandibles: Possible ontogenetic strategies and solutions. *Anat*
425 *Res Int* doi.org/10.1155/2011/484607.
- 426 Boyde A (1964) The structure and development of mammalian enamel. PhD dissertation. University of
427 London.
- 428 Boyde A (1989) Enamel. In: *Handbook of Microscopic Anatomy, Volume 6: Teeth* (eds. Oksche A
429 and Vollrath L), pp. 309–473. Berlin and New York: Springer-Verlag.
- 430 Butler PM (1956) The ontogeny of molar pattern. *Biol Rev* 31, 30–70.
- 431 Butler PM (1971) Growth of human tooth germs. In: *Dental Morphology and Evolution* (ed. Dahlberg
432 AA), pp. 3–14. Chicago: University of Chicago Press.
- 433 Cai J, Cho SW, Kim JY, Lee MJ, Cha YG, Jung HS (2007) Patterning the size and number of tooth and
434 its cusps. *Dev Biol* 304, 499–507.
- 435 Carroll SB (2008) Evo-devo and expanding evolutionary synthesis: A genetic theory of morphological

- 436 evolution. *Cell* 134, 25–36.
- 437 Chernoff B, Magwene PM (1999) Morphological Integration: Forty Years Later. In: *Morphological*
- 438 *Integration* (eds. Olsen EC, Miller RL), pp. 319–353. Chicago: University of Chicago Press.
- 439 Cheverud JM (1996) Developmental integration and the evolution of pleiotropy. *Am Zool* 36, 44–50.
- 440 Collard M, Wood B (2000) How reliable are human phylogenetic hypotheses? *Proc Natl Acad Sci U S A*
- 441 97, 5003–5006.
- 442 Collyer ML, Adams DC (2007) Analysis of two-state multivariate phenotypic change in ecological
- 443 studies. *Ecology* 88, 683–692.
- 444 Corruccini RS (1998) The dentino-enamel junction in primate mandibular molars. In: *Human Dental*
- 445 *Development, Morphology, and Pathology: A Tribute to Albert A. Dahlberg* (ed. Lukacs JR), pp. 1–16.
- 446 Portland: University of Oregon Anthropological Papers.
- 447 Dahlberg AA (1945) The changing dentition of man. *J Am Dent Assoc* 32, 676–690.
- 448 Dean MC (2000) Progress in understanding hominoid dental development. *J Anat* 197, 77–101.
- 449 Escoufier Y (1973) Le traitement des variables vectorielles. *Biometrics* 29, 751–760.
- 450 Finarelli JA, Clyde WC (2004) Reassessing hominoid phylogeny: evaluating congruence in the
- 451 morphological and temporal data. *Paleobiol* 30, 614–651.
- 452 Fincham AG, Luo W, Moradian-Oldak J, Paine ML, Snead ML, Zeichner-David M (2000) Enamel
- 453 biomineralization: the assembly and disassembly of the protein extracellular organic matrix. In:
- 454 *Development, Function and Evolution of Teeth* (eds. Teaford MF, Meredith-Smith M, Ferguson MWJ),
- 455 pp. 37–61. Cambridge: Cambridge University Press.
- 456 Grabowski MW (2013) Hominin obstetrics and the evolution of constraints. *Evol Biol* 40, 57–75.

- 457 Grine FE (2005) Enamel thickness of deciduous and permanent molars in modern *Homo sapiens*. *Am J*
458 *Phys Anthropol* 126, 14–31.
- 459 Gunz P, Mitteroecker P, Bookstein FL (2005) Semilandmarks in three dimensions. In: *Modern*
460 *Morphometrics in Physical Anthropology* (ed. Slice DE), pp. 73–98. New York: Kluwer
461 Academic/Plenum Publishers.
- 462 Hallgrímsson B, Willmore K, Hall BK (2002) Canalization, developmental stability, and morphological
463 integration in primate limbs. *Am J Phys Anthropol* 35, 131–158.
- 464 Hallgrímsson B, Jamniczky H, Young NM, et al. (2009) Deciphering the palimpsest: Studying the
465 relationship between morphological integration and phenotypic covariation. *Evol Biol* 36, 355–376.
- 466 Hansen TF (2003) Is modularity necessary for evolvability? Remarks on the relationship between
467 pleiotropy and evolvability. *Biosystems* 69, 83–94.
- 468 Harjunmaa E, Kallonen A, Voutilainen M, Hämäläinen K, Mikkola ML, Jernvall J (2012) On the
469 difficulty of increasing dental complexity. *Nature* 483, 324–327.
- 470 Harris EF, Dinh DP (2006) Intercusp relationships of the permanent maxillary first and second molars
471 in American whites. *Am J Phys Anthropol* 130, 514–28.
- 472 Hlusko LJ (2004) Integrating the genotype and phenotype in hominid paleontology. *Proc Natl Acad Sci*
473 *U S A* 101, 2653–2657.
- 474 Hlusko LJ, Suwa G, Kono R, Mahaney MC (2004) Genetics and the evolution of primate enamel
475 thickness: A baboon model. *Am J Phys Anthropol* 124, 223–233.
- 476 Hunter JP, Jernvall J (1995) The hypocone as a key innovation in mammalian evolution. *Proc Natl*
477 *Acad Sci U S A* 92, 10718–10722.

- 478 Jernvall J, Jung HS (2000) Genotype, phenotype, and developmental biology of molar tooth characters.
479 *Year Phys Anthropol* 43, 171–190.
- 480 Kavanagh KD, Evans AR, Jernvall J (2007) Predicting evolutionary patterns of mammalian teeth from
481 development. *Nature* 449, 427–432.
- 482 Klingenberg CP (2008) Morphological integration and developmental modularity. *Annu Rev Ecol Evol*
483 *Syst* 39, 115–132.
- 484 Klingenberg CP (2011) MorphoJ: an integrated software package for geometric morphometrics. *Molec*
485 *Ecol Res* 11, 353–357.
- 486 Korenhof CAW (1960) Morphogenetical Aspects of the Human Upper Molar. Utrecht:
487 Uitgeversmaatschappij Neerlandia.
- 488 Korenhof CAW (1961) The enamel-dentine border: a new morphological factor in the study of the
489 (human) molar pattern. *Proc Koninkl Nederl Acad Wetensch* 64B, 639–664.
- 490 Korenhof CAW (1982) Evolutionary trends of the inner enamel anatomy of deciduous molars from
491 Sangiran (Java, Indonesia). In: *Teeth: Form, Function and Evolution* (ed. Kurtén B), pp. 350–365. New
492 York: Columbia University Press.
- 493 Kraus BS (1952) Morphologic relationships between enamel and dentin surfaces of lower first molar
494 teeth. *J Dent Res* 31, 248–256.
- 495 Kraus BS, Jordan RE (1965) The Human Dentition Before Birth. Philadelphia: Lea and Febiger.
- 496 Lande R (1979) Quantitative genetic analysis of multivariate evolution, applied to brain: Body size
497 allometry. *Evolution* 33, 402–416.
- 498 Lande R, Arnold SJ (1983) The measurement of selection on correlated characters. *Evolution* 37, 1210–

- 499 1226.
- 500 Lewton KL (2012) Evolvability of the primate pelvic girdle. *Evol Biol* 39, 126–139.
- 501 Liversidge HM, Molleson T (2004) Variation in crown and root formation and eruption of human
- 502 deciduous teeth. *Am J Phys Anthropol* 123, 172–180.
- 503 Manly BFJ (1997) Randomization, bootstrap and Monte Carlo methods in biology. London: Chapman
- 504 & Hall.
- 505 Matsumura H, Domett KM, O'reilly DJW (2011) On the origin of pre-Angkorian peoples: perspectives
- 506 from cranial and dental affinity of the human remains from Iron Age Phum Snay, Cambodia. *Anthropol*
- 507 *Sci* 119, 67–79.
- 508 Maynard Smith J, Burian R, Kauffman S, et al. (1985) Developmental constraints and evolution. *Q Rev*
- 509 *Biol* 60, 265–287.
- 510 Marroig G, Shirai L, Porto A, de Oliveira F, De Conto V (2009) The evolution of modularity in the
- 511 mammalian skull II: Evolutionary consequences. *Evol Biol* 36, 136–148.
- 512 Miller GS (1918) The Piltdown jaw. *Am J Phys Anthropol* 1, 25–52.
- 513 Nager G, (1960) Der vergleich zwischen dem räumlichen verhalten des dentinkronenreliefs und dem
- 514 schmelzrelief der zahnkrone. *Acta Anat* 42, 226–250.
- 515 Nanci A (2013) Ten Cate's Oral Histology: Development, Structure and Function, 8th Edition. St.
- 516 Louis: Mosby Elsevier.
- 517 Olejniczak AJ, Gilbert CG, Martin LB, Smith TM, Ulhaas L, Grine FE (2007) Morphology of the
- 518 enamel-dentine junction in sections of anthropoid primate maxillary molars. *J Hum Evol* 53, 292–301.
- 519 Ortiz A, Skinner MM, Bailey SE, Hublin JJ (2012) Carabelli's trait revisited: an examination of

- 520 mesiolingual features at the enamel-dentine junction and enamel surface of Pan and Homo sapiens
- 521 upper molars. *J Hum Evol* 63, 586–596.
- 522 Pilbrow V (2006) Population systematics of chimpanzees using molar morphometrics. *J Hum Evol* 51,
- 523 646–662.
- 524 Porto A, de Oliveira FB, Shirai LT, de Conto V, Marroig G (2009) The evolution of modularity in the
- 525 mammalian skull I: Morphological integration patterns and magnitudes. *Evol Biol* 36, 118–135.
- 526 R Development Core Team (2011) R: A Language and Environment for Statistical Computing. Vienna:
- 527 R Foundation for Statistical Computing. <http://cran.R-project.org>.
- 528 Rohlf FJ, Slice D (1990) Extensions of the Procrustes method for the optimal superimposition of
- 529 landmarks. *Syst Zool* 39, 40–59.
- 530 Sakai T, Hanamura H (1971) A morphology study of enamel-dentin border on the Japanese dentition.
- 531 Part V. Maxillary molar. *J Anthropol Soc Nippon* 79, 297–322.
- 532 Sakai T, Hanamura H (1973) A morphology study of enamel-dentin border on the Japanese dentition.
- 533 Part VII. General conclusion. *J Anthropol Soc Nippon* 81, 87–102.
- 534 Sasaki K, Kanazawa E (1999) Morphological traits on the dentino-enamel junction of lower deciduous
- 535 molar series. In: *Dental morphology 1998: Proceedings of the 11th international symposium on dental*
- 536 *morphology* (eds. Mayhall J, Heikkinen T), pp. 167–178. Oulu: Oulu University Press.
- 537 Saunders SR, Mayhall JT (1982) Developmental patterns of human morphological traits. *Archs Oral*
- 538 *Biol* 27, 45–49.
- 539 Shellis RP (1984) variations in growth of the enamel crown in human teeth and a possible relationship
- 540 between growth and enamel structure. *Archs Oral Biol* 29, 697–705.

- 541 Simmer JP, Papagerakis P, Smith CE, et al. (2010) Regulation of dental enamel shape and hardness. *J*
- 542 *Dent Res* 89, 1024–1038.
- 543 Simons EL, Pilbeam D (1972) Hominoid paleoprimatology. In: *The Functional and Evolutionary*
- 544 *Biology of Primates* (ed. Tuttle R), pp. 36–62. Chicago: Aldine-Atherton.
- 545 Skinner MM, Wood BA, Boesch C, et al. (2008) Dental trait expression at the enamel-dentine junction
- 546 of lower molars in extant and fossil hominoids. *J Hum Evol* 54, 173–186.
- 547 Skinner MM, Wood BA, Hublin JJ (2009) Protostylid expression at the enameledentine junction and
- 548 enamel surface of mandibular molars of *Paranthropus robustus* and *Australopithecus africanus*. *J Hum*
- 549 *Evol* 56, 76–85.
- 550 Skinner MM, Evans A, Smith T, et al. (2010) Brief Communication: Contributions of enamel-dentine
- 551 junction shape and enamel deposition to primate molar crown complexity. *Am J Phys Anthropol* 142,
- 552 157–163.
- 553 Skinner MM, Gunz P (2010) The presence of accessory cusps in chimpanzee lower molars is consistent
- 554 with a patterning cascade model of development. *J Anat* 217, 245–253.
- 555 Smith P, Gomorri JM, Spitz S, Becker J (1997) Model for the examination of evolutionary trends in
- 556 tooth development. *Am J Phys Anthropol* 102, 283–294.
- 557 Smith P, Gomori JM, Shaked R, Haydenblit R, Joskowicz L (2000) A computerized approach to
- 558 reconstruction of growth patterns in hominid molar teeth. In: *Proceedings of the 11th International*
- 559 *Symposium on Dental Morphology* (eds. Mayhall J, Heikkinen T), pp. 388–397. Oulu: Oulu University
- 560 Press.
- 561 Smith P, Avishai G, Muller R, Gabet Y (2011) Computerized reconstruction of prenatal growth

- 562 trajectories in the dentition: Implications for the taxonomic status of Neandertals. In: *Continuity and*
563 *discontinuity in the peopling of Europe: One hundred fifty years of Neanderthal study* (eds. Condemi S,
564 Weniger G-C), pp. 165–173. New York: Springer Science+Business Media B.V.
- 565 Sokal RR, Braumann CA (1980) Significance tests for coefficients of variation and variability profiles.
566 *Syst Zool* 29, 50–66.
- 567 Suzuki M, Sakai T (1973) Occlusal surface pattern of the lower molars and the second deciduous molar
568 among living Polynesians. *Am J Phys Anthropol* 39, 305–315.
- 569 Townsend G, Richards L, Hughes T (2003) Molar intercuspal dimensions: genetic input to phenotypic
570 variation. *J Dent Res* 82, 350–355.
- 571 Villmoare B, Fish J, Jungers W (2011) Selection, morphological integration, and strepsirrhine
572 locomotor adaptations. *Evol Biol* 38, 88–99.
- 573 Wagner GP (1984) On the eigenvalue distribution of genetic and phenotypic dispersion matrices:
574 Evidence for a nonrandom organization of quantitative character variation. *J Math Biol* 21, 77–95.
- 575 Wagner GP, Altenberg L (1996) Complex adaptations and the evolution of evolvability. *Evolution* 50,
576 967–976.
- 577 Wagner GP, Booth G, Bagheri-Chaichian H (1997) A population genetic theory of canalization.
578 *Evolution* 51, 329–347.
- 579 Willmore KE, Young N, Richtsmeier JT (2007) Phenotypic variability: its components, measurement
580 and underlying developmental processes. *Evol Biol* 34, 99–120.

581 **Tables**

Table 1 Results of PLS analyses between EDJ and OES corresponding to UM1 and um2

	UM1			um2		
	% Total Cov.	Correlation coefficient	P-value ¹	% Total Cov.	Correlation coefficient	P-value ¹
1	49.43	0.951	<0.001	43.14	0.974	<0.001
2	17.39	0.933	<0.001	25.11	0.970	<0.001
3	14.65	0.908	<0.001	17.76	0.954	<0.001
4	10.22	0.879	<0.001	6.52	0.948	<0.001

¹Randomization rounds: 1000

582

Table 2 Results of principal component analysis with the total sample

	Eigenvalue	% Explained variance	% Cumulative variance
1	0.0016	19.99	19.99
2	0.0012	14.86	34.85
3	0.0009	11.80	46.64
4	0.0007	9.08	55.73
5	0.0005	6.86	62.58
6	0.0005	6.68	69.26
7	0.0004	5.31	74.58
8	0.0002	3.14	77.71
9	0.0002	2.90	80.61
10	0.0002	2.35	82.96

583



Figure legends





Figure 1. Digital image of permanent maxillary first molar crown (lingual view). (a) EDJ ridge curve digitized on the EDJ surface. (b) OES ridge curve digitized on the OES. Red circles are landmarks, and yellow circles are semi-landmarks. Numbers appended to each section of the ridge curve refer to the equally-spaced interpolated semi-landmarks.

Figure 2. Scatter plots representing the first and second pairs of PLS axes between EDJ and OES within the same tooth class. (a) PLS1 UM1, (b) PLS2 UM1, (c) PLS1 um2, (d) PLS2 um2. Shape deformation corresponding to each axis is provided to the left of x-axes or above y-axes. Each shape deformation is represented in colored line whose scale factor used for is 0.1 and mean shape is represented in gray line.


Figure 3. Principal component plots for shape variation between EDJ and OES of both UM1 and um2. (a) Plots of PC1 versus PC2 scores. Variance explained by PC1 and PC2 is 34.85% of total variance. Shape deformation corresponding to the positive or negative loadings of each axis is provided to the left and right for x-axes or the above and bottom for y-axes. Each shape deformation is represented in colored line whose scale factor used for is 0.1 and mean shape is represented in gray line. Arrows show morphological change vectors from mean shape represented in large symbols of EDJ to that of OES for each tooth class. (b) Relationship between EDJ and OES for PC1 in both UM1 and um2. The slope and intercept of the regression line for UM1 are 0.804 and -0.070, respectively ($r=0.51$, $P<0.001$). The slope and intercept of the regression line for um2 are 0.876 and -0.068, respectively ($r=0.84$, $P<0.001$). (c) Relationship between EDJ and OES for PC2 in both UM1 and um2. The slope and intercept of the

605 regression line for UM1 are 0.863 and -0.002, respectively ($r=0.92$, $P<0.001$). The slope and intercept of
606 the regression line for um2 are 0.918 and 0.007, respectively ($r=0.95$, $P<0.001$).
607
608 Figure 4. (a) Bar graph showing the size variation for four configurations (UM1EDJ, UM1OES,
609 um2EDJ and um2OES). Significance test for coefficient of variation for LogCS among them reveals
610 that there is no significant difference ($P>0.05$). (b) Bar graph showing mean of propcrustes distance
611 from each mean shape for shape variance of four configurations (UM1EDJ, UM1OES, um2EDJ and
612 um2OES), and the error bars show standard deviations. The Kruskal-Wallis test reveals a significant
613 difference among them ($P<0.001$). A nonparametric multiple-comparison test between EDJ and OES
614 within the same tooth class reveals that the difference is highly significant in UM1 ($P<0.001$). (c) Bar
615 graph showing the scaled variances of eigenvalue for morphological integration for four configurations
616 (UM1EDJ, UM1OES, um2EDJ and um2OES). The error bars shown are standard deviations obtained
617 by resampling the original datasets with replacement 1000 times. Bootstrap tests between EDJ and OES
618 within the same tooth class reveal that the difference is highly significant only in UM1 ($P=0.009$). (d)
619 Bar graph showing the evolutionary flexibility for four configurations (UM1EDJ, UM1OES, um2EDJ
620 and um2OES). The error bars shown are standard deviations obtained by resampling the original
621 datasets with replacement 1000 times. Bootstrap tests between EDJ and OES within the same tooth
622 class reveal that there is no significant difference ($P>0.05$).






a



b



c

